

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE January 25, 1996	3. REPORT TYPE AND DATES COVERED preprint of journal article
4. TITLE AND SUBTITLE Molecular Electrostatic Potentials as Indicators of Covalent Radii			5. FUNDING NUMBERS N00014-95-1-0028 Dr. Richard S. Miller R&T Code 33e 1806
6. AUTHOR(S) John J. M. Wiener, M. Edward Grice, Jane S. Murray and Peter Politzer			8. PERFORMING ORGANIZATION REPORT NUMBER 88
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of New Orleans Department of Chemistry New Orleans, Louisiana 70148			10. SPONSORING/MONITORING AGENCY REPORT NUMBER
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research Code 333 800 N. Quincy Street Arlington, VA 22217			
11. SUPPLEMENTARY NOTES			
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release. Unlimited distribution.		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Through nonlocal density functional calculations (B/LYP/6 -31G**), we show that the minimum of the molecular electrostatic potential along an internuclear axis provides a more realistic boundary point between two bonded atoms than does the electronic density minimum. We also suggest that the two bonded atoms contribute approximately equally to the magnitude of the minimum potential along the axis. <div style="text-align: right;">DTIC QUALITY INSPECTED 1</div>			
14. SUBJECT TERMS nonlocal density functional theory; electronic density; electrostatic potential; covalent radii			15. NUMBER OF PAGES 9
			16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited

19960208 080

OFFICE OF NAVAL RESEARCH

CONTRACT N00014-95-1-0028

R&T Code 4131D02

Dr. Richard S. Miller

Technical Report No. 88

Molecular Electrostatic Potentials as Indictors of Covalent Radii

by

John J. M. Wiener, M. Edward Grice, Jane S. Murray and Peter Politzer

Prepared for Publication

in

Journal of Chemical Physics

Department of Chemistry
University of New Orleans
New Orleans, LA 70148

January 25, 1996

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

The electrostatic potential $V^{\text{at}}(\mathbf{r})$ in the space around an atom with nuclear charge Z and electronic density $\rho^{\text{at}}(\mathbf{r})$ is given by eq. (1), which invokes the spherical symmetry of the charge distribution.¹

$$V^{\text{at}}(\mathbf{r}) = V^{\text{at}}(r) = \frac{Z}{r} - \int \frac{\rho^{\text{at}}(\mathbf{r}') d\mathbf{r}'}{|\mathbf{r}' - \mathbf{r}|} \quad (1)$$

For neutral atoms, $V^{\text{at}}(r)$ decreases monotonically with radial distance from the nucleus.² Figure 1 shows that $\ln V^{\text{at}}(r)$ varies nearly linearly with radial distance in the valence region, indicating that the relationship is essentially exponential, $V^{\text{at}}(r) = A \exp(-Br)$. In Figure 1 are given $\ln V^{\text{at}}(r)$ equations for nine atoms, obtained by fitting the electrostatic potentials computed from Hartree-Fock atomic wave functions.³ These equations, all of which have correlation coefficients better than 0.997, cover the ranges $r_0 \leq r \leq 3$ bohr, where r_0 was taken to be the radius of the inner boundary of the valence shell, defined as the radial distance to the outermost inflection point of the average local ionization energy of the atom.⁴ (Other criteria could also be used for defining this boundary,⁵⁻⁷ without significantly affecting our conclusions.)

The electrostatic potential of a molecule with electronic density $\rho^{\text{mol}}(\mathbf{r})$ is given by eq. (2):

$$V^{\text{mol}}(\mathbf{r}) = \sum_A \frac{Z_A}{|\mathbf{R}_A - \mathbf{r}|} - \int \frac{\rho^{\text{mol}}(\mathbf{r}') d\mathbf{r}'}{|\mathbf{r}' - \mathbf{r}|} \quad (2)$$

Z_A is the charge on nucleus A , located at \mathbf{R}_A . Pathak and Gadre have shown that a molecular electrostatic potential has no local maxima.⁸ (At nuclear sites, it is of course undefined.) Along the axis between two bonded atoms, therefore, $V^{\text{mol}}(\mathbf{r})$ must decrease monotonically with distance from each nucleus and reach an axial minimum at some point \mathbf{r}_m .⁹ Since $(\partial V^{\text{mol}} / \partial r_{\text{axial}})_{\mathbf{r}_m} = 0$, an infinitesimal charge δq placed at \mathbf{r}_m will feel no electrostatic force from either direction along the internuclear axis. We have accordingly explored the possibility that \mathbf{r}_m may be a meaningful axial boundary point between the two bonded atoms.

We have computed optimized geometries, electronic densities and electrostatic potentials for the 19 molecules listed in Table I. The calculations were carried out using a non-local density functional procedure (Gaussian 94,¹⁰ Becke exchange¹¹ and Lee, Yang and Parr correlation functionals,¹² 6-31G** basis set). For the bonds indicated, we found the distances from each nucleus to the minima of both $V_{\text{mol}}(\mathbf{r})$ and (following Bader *et al*¹³) $\rho^{\text{mol}}(\mathbf{r})$ along the internuclear axis. As a test to determine whether either of these points is an effective boundary indicator, we compare these distances to literature values of the covalent radii of the respective atoms.

Overall, the distances to the minima of $V_{\text{mol}}(\mathbf{r})$ are in reasonable agreement with the covalent radii, especially in view of the fact that the latter do not refer to any specific molecule but

rather were obtained from experimental structural studies of a number of compounds.¹⁴ The greatest deviations occur for some of the hydrogens. On the other hand, the distances to the minima of $\rho^{\text{mol}}(\mathbf{r})$ usually differ significantly more from the covalent radii and show some major discrepancies, more than 0.30 Å in ten instances. The relative radii given by the density minima are often not what would be anticipated; particularly striking examples are PH_3 and SiH_4 , in which the hydrogens are larger than the central atoms. The electron density wells also tend to be wider and flatter than those of the electrostatic potential. Table II summarizes the data in Table I, giving for each atom the averages of the covalent radii obtained from the potential minima and the density minima. The former are closer to the literature values, often considerably so, for every atom.

These results indicate that the location of the minimum electrostatic potential along the internuclear axis does define a meaningful boundary between two bonded atoms. The less realistic covalent radii obtained from the minimum electronic density are consistent with the unexpectedly large magnitudes of the atomic charges found by using this point on the internuclear axis as the basis for partitioning the electronic charge.^{13,15} (For a relevant critical analysis, see Perrin.¹⁶)

If $V^{\text{mol}}(\mathbf{r})$ at its minimum on the internuclear axis, i.e. $V^{\text{mol}}(\mathbf{r}_m)$, can be regarded as reflecting largely the electrostatic potentials of the two bonded atoms A and B, then within this approximation, the point \mathbf{r}_m corresponds to the intersection of $V^A(\mathbf{r})$ and $V^B(\mathbf{r})$, and their contributions to $V^{\text{mol}}(\mathbf{r}_m)$ should be equal. (This does not preclude other contributions to $V^{\text{mol}}(\mathbf{r}_m)$.) One can arrive at the same conclusion by noting that several proposed definitions of electronegativity express it as an electrostatic potential or force created by an atom at a distance equal to its covalent radius.¹⁷⁻¹⁹ It then follows from the principle of electronegativity equalization²⁰⁻²² that these potentials or forces should be the same for two bonded atoms at their internuclear axial boundary point.

To test the proposed condition $V^A(\mathbf{r}_m) \approx V^B(\mathbf{r}_m)$, we have found the points of intersection of all of the various pairwise combinations of the $\ln V(\mathbf{r})$ equations in Figure 1, when the respective nuclei are separated by the sum of the literature covalent radii of the atoms. The objective was to determine how well the radial distances to the points of intersection reproduce the literature covalent radii.

For those combinations in which both atoms are in the first row of the periodic table (C - F) or both are in the second row (Si - Cl), the differences between the calculated and the literature covalent radii are very small, the average being 0.013 Å. Excellent agreement is also obtained for the second row/third row pairs (Si-Br, P-Br, S-Br and Cl-Br); the average discrepancy is 0.015 Å. The deviations are larger for combinations of a first-row atom with one in the second row or with bromine; the average difference is 0.066 Å. Among these, the covalent radius of the first-row atom is invariably overestimated and that of the second-row atom or bromine is underestimated.

This may reflect the well-known discontinuities in many properties between atoms of the first and subsequent rows.²³⁻²⁷ Overall, these results do provide significant support for the idea that bonded atoms contribute approximately equally to $V^{\text{mol}}(\mathbf{r}_m)$.

The main point that we have tried to make in this paper is that the electrostatic potential minimum along the internuclear axis provides a realistic boundary point between two bonded atoms, more so than does the electronic density minimum. (In this context it is relevant to recall earlier work indicating that the locations of the electrostatic potential minima of monoatomic negative ions can be identified with their ionic radii.²⁸) We have also suggested that the two bonded atoms contribute approximately equally to the minimum potential along the axis. It is reasonable to suppose that these conclusions are related to the observation that the covalent radius of an atom is quite close to the radial distance at which its electrostatic potential equals its chemical potential.²⁹ The relationship is being investigated.

Acknowledgment:

We greatly appreciate the financial support of the Office of Naval Research, through contract No. N00014-95-1-0028 and Program Officer Dr. Richard S. Miller.

Figure Caption:

Figure 1. Natural logarithm of calculated electrostatic potential, in kcal/mole, plotted against radial distance, in bohr, for the atoms indicated. The equations describing these plots in the valence regions of the atoms are also given. (See text for definition of valence region.)

References

- 1 G. Delgado-Barrio and R. F. Prat, Phys. Rev. A **12**, 2288 (1975).
- 2 H. Weinstein, P. Politzer, and S. Srebrenik, Theor. Chim. Acta **38**, 159 (1975).
- 3 E. Clementi, *Tables of Atomic Functions* (IBM, San Jose, CA, 1965).
- 4 P. Politzer, J. S. Murray, M. E. Grice, T. Brinck, and S. Ranganathan, J. Chem. Phys. **95**, 6699 (1991).
- 5 P. Politzer and R. G. Parr, J. Chem. Phys. **64**, 4634 (1976).
- 6 K. D. Sen, M. Slamet, and V. Sahni, Chem. Phys. Lett. **205**, 313 (1993).
- 7 M. Kakkar and K. D. Sen, Chem. Phys. Lett. **226**, 241 (1994).
- 8 R. K. Pathak and S. R. Gadre, J. Chem. Phys. **93**, 1770 (1990).
- 9 S. R. Gadre, S. A. Kulkarni, and I. H. Shrivastava, J. Chem. Phys. **96**, 5253 (1992).

- 10 M. J. Frisch, G. W. Trucks, H. B. Schlegel, P. M. W. Gill, B. G. Johnson, M. A. Robb, J. R. Cheeseman, T. A. Keith, G. A. Petersson, J. A. Montgomery, K. Raghavachari, M. A. Al-Laham, V. G. Zakrzewski, J. V. Ortiz, J. B. Foresman, J. Cioslowski, B. B. Stefanov, A. Nanayakkara, M. Challacombe, C. Y. Peng, P. Y. Ayala, W. Chen, M. W. Wong, J. L. Andres, E. S. Replogle, R. Gomperts, R. L. Martin, D. J. Fox, J. S. Binkley, D. J. Defrees, J. Baker, J. P. Stewart, M. Head-Gordon, C. Gonzalez, and J. A. Pople, GAUSSIAN 94 (Revision B3) (Gaussian, Inc., Pittsburgh, PA, 1995).
- 11 A. D. Becke, Phys. Rev. A **38**, 3098 (1988).
- 12 C. Lee, W. Yang, and R. G. Parr, Phys. Rev. B **37**, 785 (1988).
- 13 R. F. W. Bader, P. M. Beddall, and P. E. Cade, J. Am. Chem. Soc. **93**, 3095 (1971).
- 14 L. Pauling, *The Nature of the Chemical Bond*, 3rd. ed. (Cornell University Press, Ithaca, NY, 1960).
- 15 R. F. W. Bader, Acc. Chem. Res. **8**, 34 (1975); **18**, 9 (1985).
- 16 C. L. Perrin, J. Am. Chem. Soc. **113**, 2865 (1991).
- 17 W. Gordy, Phys. Rev. **69**, 604 (1946).
- 18 A. L. Allred and E. G. Rochow, J. Inorg. Nucl. Chem. **5**, 264, 269 (1958).
- 19 R. J. Boyd and K. E. Edgecombe, J. Am. Chem. Soc. **110**, 4182 (1988).
- 20 R. T. Sanderson, J. Am. Chem. Soc. **74**, 272 (1952); Science **121**, 207 (1955).
- 21 R. G. Parr, R. A. Donnelly, M. Levy, and W. E. Palke, J. Chem. Phys. **68**, 3801; R. A. Donnelly and R. G. Parr, J. Chem. Phys. **69**, 4431 (1978) (1978).
- 22 P. Politzer and H. Weinstein, J. Chem. Phys. **71**, 4218 (1979).
- 23 R. T. Sanderson, *Inorganic Chemistry* (Reinhold, New York, 1967) pp. 285-292.
- 24 P. Politzer, in *Homoatomic Rings, Chains and Macromolecules of Main-Group Elements*, edited by A. L. Rheingold (Elsevier, New York, 1977), Chapter 4.
- 25 F. A. Cotton and G. Wilkinson, *Inorganic Chemistry* (John Wiley and Sons, New York, 1980), pp. 208-211.
- 26 B. E. Douglas, D. H. McDaniel, and J. Alexander, *Concepts and Models of Inorganic Chemistry*, 2nd ed. (Wiley, New York, 1983) pp. 45-47, 226.
- 27 P. Politzer, J. E. Huheey, J. S. Murray, and M. Grodzicki, J. Mol. Struct. (Theochem) **259**, 99 (1992).
- 28 K. D. Sen and P. Politzer, J. Chem. Phys. **90**, 4370 (1989); **91**, 5123 (1989).
- 29 P. Politzer, R. G. Parr, and D. R. Murphy, J. Chem. Phys. **79**, 3859 (1983); Phys. Rev. B **31**, 6809 (1985).

TABLE I. Literature values of covalent radii compared to results obtained in terms of axial minima of (a) electrostatic potential and (b) electronic density.

Molecule	Bond, A-B	Bond Length, Å			Covalent radii, Å		
		Calc.	Exp. ^a		Literature ^b	Min., V _{mol}	Min., ρ ^{mol}
CH ₄	C-H	1.099	1.094	C: H:	0.772 0.30	0.699 0.400	0.686 0.413
HCN	C-H	1.076	1.065	C: H:	0.772 0.30	0.683 0.393	0.703 0.373
NH ₃	N-H	1.028	1.024	N: H:	0.70 0.30	0.656 0.372	0.754 0.274
H ₂ O	O-H	0.976	0.972	O: H:	0.66 0.30	0.623 0.353	0.774 0.202
SiH ₄	Si-H	1.494	1.481	Si: H:	1.17 0.30	1.01 0.487	0.724 0.769
PH ₃	P-H	1.436	1.427	P: H:	1.10 0.30	0.971 0.465	0.699 0.737
H ₂ S	S-H	1.359	1.336	S: H:	1.04 0.30	0.921 0.438	0.837 0.521
CH ₃ NH ₂	C-N	1.480	1.471	C: N:	0.772 0.70	0.748 0.732	0.620 0.860
CH ₃ NO ₂	C-N	1.520	1.489	C: N:	0.772 0.70	0.768 0.752	0.617 0.903
(H ₃ C) ₂ O	C-O	1.427	1.410	C: O:	0.772 0.66	0.729 0.698	0.512 0.914
C ₂ H ₅ OH	C-O	1.441	1.431	C: O:	0.772 0.66	0.738 0.703	0.523 0.918
H ₃ C-OCi	C-O	1.439	1.418	C: O:	0.772 0.66	0.735 0.703	0.513 0.925
CH ₃ F	C-F	1.398	1.383	C: F:	0.772 0.64	0.724 0.674	0.482 0.916
CH ₃ PH ₂	C-P	1.896	1.863	C: P:	0.772 1.10	0.846 1.05	1.14 0.754
CH ₃ SH	C-S	1.857	1.819	C: S:	0.772 1.04	0.836 1.02	0.875 0.983
CH ₂ Cl ₂	C-Cl	1.814	1.772	C: Cl:	0.772 0.99	0.829 0.985	0.793 1.02

(Continued)

TABLE I. Literature values of covalent radii compared to results obtained in terms of axial minima of (a) electrostatic potential and (b) electronic density (continued).

Molecule	Bond, A-B	Bond Length, Å			Covalent radii, Å		
		Calc.	Exp. ^a		Literature ^b	Min., V_{mol}	Min., ρ_{mol}
FNNF	F-N	1.428	1.385	F:	0.64	0.691	0.778
				N:	0.70	0.737	0.648
FSSF	F-S	1.698	1.635	F:	0.64	0.722	0.988
				S:	1.04	0.976	0.710
SiH ₃ Cl	Si-Cl	2.098	2.049	Si:	1.17	1.08	0.759
				Cl:	0.99	1.02	1.34

^aM. D. Harmony, V. W. Laurie, R. L. Kuczkowski, R. H. Schwendeman, D. A. Ramsay, F. J. Lovas, W. J. Lafferty, and A. G. Maki, *J. Phys. Chem. Ref. Data* **8**, 619 (1979).

^bJ. A. Dean, *Lange's Handbook of Chemistry*, 14th ed. (McGraw-Hill, New York, 1992).

TABLE II. Literature covalent radii compared to averages of values given in Table I, correspondingly to axial minima of (a) electrostatic potential and (b) electronic density.

Atom	Number of values	Covalent radii, Å		
		Literature	Average, min., V^{mol}	Average, min., ρ^{mol}
H	7	0.30	0.415	0.470
C	11	0.772	0.758	0.679
N	4	0.70	0.719	0.791
O	4	0.66	0.682	0.883
F	3	0.64	0.696	0.894
Si	2	1.17	1.05	0.742
P	2	1.10	1.01	0.727
S	3	1.04	0.97	0.843
Cl	2	0.99	1.00	1.18

